

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM
maintained by the
CALIFORNIA DEPARTMENT OF FISH AND GAME
and supported by the
CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP
Database Version 8.1 (2005)

M103 Pacific Kangaroo Rat *Dipodomys agilis*
Family: Heteromyidae Order: Rodentia Class: Mammalia

Written by: G. Ahlborn
Reviewed by: M. White
Edited by: M. White

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

Common, yearlong resident of southern California. Inhabits South Coast region, generally west of the Mojave and Colorado deserts, from the Mexican border north to the Santa Barbara-San Luis Obispo Co. line. Also occurs in Tehachapi and Piute Mts. inland to South Fork of Kern River in southeastern Tulare Co. Found in coastal scrub, open stages of chaparral, and desert scrub habitats, and in some conifer woodlands. It occurs in habitats below 2250 m (7400 ft).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Primary foods include seeds of grasses, forbs, and shrubs (Meserve 1976a). Forages on open ground adjacent to shrubs, searching for, and gathering, food. Large amounts of food can be held in cheek pouches, some of which later is stored in underground caches.

Cover: Burrows are excavated in friable soils. Shrubs and herbage provide cover during above-ground activities.

Reproduction: Nests in burrows excavated in friable soils.

Water: Probably needs drinking water (MacMillen 1964).

Pattern: Open areas with scattered vegetation and friable soils provide suitable habitat for *D. agilis*.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong nocturnal activity.

Seasonal Movements/Migration: Non-migratory.

Home Range: Home ranges varied from 0.1 to 0.6 ha (0.4 to 1.5 ac), with a mean of 0.3 ha (0.8 ac) (MacMillen 1964).

Territory: Females may be territorial during breeding season. Territory and home range probably are the same size.

Reproduction: Reproduction occurs from January into July. Litter size ranges from 2-4; average = 2.6.

Niche: *D. agilis* is closely related to *D. heermanni* and *D. venustus*. Unlike most other

Dipodomys, *D. agilis* is not well adapted to extreme xeric conditions. Rattlesnakes, owls, badgers, foxes, skunks, weasels, and coyotes are likely predators. Other seed-eating species may compete when food is scarce.

REFERENCES

- Bleich, V. C. 1973. Ecology of rodents at the United States Naval Weapon Station, Seal Beach, Fallbrook Annex, San Diego County, California. M.A. Thesis, California State Univ., Long Beach. 102pp.
- Bontrager, D. R. 1973. Rodent ecology of the Santa Rosa Plateau, Riverside County, California. M.A. Thesis, Calif. State Univ., Long Beach. 115pp.
- Christopher, E. A. 1973. Sympatric relationships of the kangaroo rats, *Dipodomys merriami* and *Dipodomys agilis*. J. Mammal. 54:317-326.
- Grinnell, J. 1922. A geographical study of the kangaroo rats of California. Univ. Calif. Publ. Zool. 24:1-124.
- MacMillen, R. E. 1964. Population ecology, water relations and social behavior of a southern California semidesert rodent fauna. Univ. Calif. Publ. Zool. 71:1-59.
- Meserve, P. L. 1976a. Habitat and resource utilization by rodents of a California coastal sage scrub community. J. Anim. Ecol. 45:647-666.
- Meserve, P. L. 1976b. Food relationships of a rodent fauna in a California coastal sage scrub community. J. Mammal. 57:300-319.
- Stock, A. D. 1974. Chromosome evolution in the genus *Dipodomys* and its taxonomic and phylogenetic implications. J. Mammal. 55:505-527.